

$B \rightarrow \pi\pi$ decays: branching ratios and CP asymmetries

A.B. Kaidalov* and M.I. Vysotsky†

Institute of Theoretical and Experimental Physics,
Moscow, Russia

Abstract

Theoretically motivated smallness of the penguin amplitude in $B \rightarrow \pi\pi$ decays allows to calculate the value of the unitarity triangle angle $\alpha(\phi_2)$ with good accuracy. The relatively large branching ratio of the decay into $\pi^0\pi^0$ is explained by the large value of FSI phase difference between decay amplitudes with $I = 0$ and $I = 2$.

PACS: 12.15.Hh, 13.20.He

1 Introduction

The exclusive decay amplitudes of hadrons are determined by dynamics at large distances and can not be calculated with presently available QCD tools.

Fortunately it was found long ago that the experimental data on branching ratios and CP asymmetries of $B \rightarrow \pi\pi$ decays allow to determine the value of the unitarity triangle angle α with essentially no hadronic input using isospin invariance of strong interactions only [1]. However, large experimental uncertainties in particular in the values of the direct CP asymmetries lead to poor accuracy in the value of α determined in this way.

If the penguin amplitudes are negligible in charmless strangeless B decays we would determine the value of unitarity triangle angle α from CP asymmetry S_{+-} extracted from $B \rightarrow \pi^+\pi^-$ decay data with essentially no theoretical uncertainties. As it was found in paper [2] neglecting penguin amplitudes one gets the values of angle α from CP asymmetries in B_d decays to $\pi^+\pi^-$, $\rho^+\rho^-$ and $\pi^\pm\rho^\mp$ consistent with the global fit of unitarity triangle. Since the penguin contributions to these decays are different [3] the fact that the numerical values of α

*kaidalov@itep.ru

†vysotsky@itep.ru

are close to each other testifies in favor of smallness of penguin amplitudes. Small penguin corrections to these decay amplitudes were accounted for in [4] where the hadronic amplitudes were found from the quark amplitudes with the help of factorization. However, it is well known that the branching ratio of $B_d(\bar{B}_d) \rightarrow \pi^0\pi^0$ decay predicted by factorization appears to be more than 10 times smaller than the experimental data. The way out of this contradiction could be large FSI phases in $B \rightarrow \pi\pi$ decays. The validity of this theoretical ingredient will be checked by the more accurate experimental data.

Though the penguin contribution is relatively small compared to tree amplitudes and can be neglected in the first approximation in the decay probabilities and in the CPV parameters S it determines the CPV parameters C and should be accounted for in the analysis of the complete set of observables.

The charmless strangeless B decays are described by $b \rightarrow u\bar{u}d$ quark transition. The effective Hamiltonian responsible for this transition consists of two parts: the tree level weak amplitude (operators O_1 and O_2 in standard notations) dressed by gluons and the gluon penguin amplitudes (operators $O_3 - O_6$); the parametrically small electroweak penguins are omitted. The gluon penguins being very important in $\Delta S = 1$ strange particles nonleptonic weak decays are almost negligible in $\Delta B = 1$, $\Delta S = 0$ transitions. The reason is twofold: firstly, Wilson coefficients are much smaller in case of B decays because infrared cutoff is at $\mu \sim m_b$ instead of $\mu \sim \Lambda_{QCD}$; secondly, the enhancement factor originated from the right-handed currents $m_\pi^2/m_s(m_u + m_d) \sim 10$ for strange particles decays is replaced by $m_\pi^2/m_b(m_u + m_d) \sim 1/3$ for beauty hadrons. That is why after presenting the general phenomenological expressions for the amplitudes we will start our analysis of $B \rightarrow \pi\pi$ decays in Section 2 by the sequestered Hamiltonian which does not contain penguin contributions¹. From the experimental data on $B_d(\bar{B}_d) \rightarrow \pi^+\pi^-, \pi^0\pi^0$ and $B_u \rightarrow \pi^+\pi^0$ branching ratios we will extract the moduli of the amplitudes of the decays into $\pi\pi$ states with isospin zero A_0 and two A_2 and find the final state interaction (FSI) phase shift $\delta \equiv \delta_2 - \delta_0$ between these two amplitudes. The value of the unitarity triangle angle α in this approximation is directly determined by CP asymmetry S_{+-} .

¹Let us stress that while from the smallness of $B \rightarrow \pi^0\pi^0$ decay width it would follow that penguins are small, the opposite statement is not correct: the relatively large width to neutral pions does not necessary mean that penguins are large.

While the absolute values of the amplitudes A_0 and A_2 are reproduced with good accuracy by the factorization formulas, the FSI phase shift appears to be unexpectedly large, $\delta = -(53^\circ \pm 7^\circ)$. This is the reason why $B \rightarrow \pi^0 \pi^0$ decay probability is significantly enhanced in comparison with the naive factorization approach, where one neglects δ . In Section 3 FSI phase differences in $K \rightarrow \pi\pi$, $D \rightarrow \pi\pi$ and $B \rightarrow D\pi$ decays are considered. In all these cases the phases are large, which is attributed to the existence of $I = 0$ resonances in $\pi\pi$ scattering in the cases of $K \rightarrow \pi\pi$ and (partly) in $D \rightarrow \pi\pi$ decays while large FSI phases in $B \rightarrow D\pi$ decays are unexpected. In Section 4 we consider the theoretical estimates of δ and show how FSI can enhance B width to neutral pions not enhancing that to neutral ρ mesons in accordance with experimentally observed suppression of $B \rightarrow \rho^0 \rho^0$ decay width.

In Section 5 the penguin contributions are considered; the corrections to the numerical values of A_0 and δ due to gluon penguin amplitudes are determined, as well as the correction to the unitarity triangle angle α and the values of CP asymmetries C_{+-} and C_{00} . In Conclusions the pattern of the $B \rightarrow \pi\pi$ decay amplitudes emerging from the experimental data is presented. Appendix contains the calculations of the decay amplitudes in factorization approximation.

2 $B \rightarrow \pi\pi$ without penguins: decay amplitudes from branching ratios

The quark Hamiltonian responsible for $B \rightarrow \pi\pi$ decays has the parts with $\Delta I = 1/2$ and $\Delta I = 3/2$ which produce π -mesons in the states with $I = 0$ and $I = 2$ correspondingly. QCD penguins having $\Delta I = 1/2$ contribute only to the $I = 0$ amplitude. Taking into account the corresponding Clebsch–Gordan coefficients and separating the penguin contribution (P) with the CKM phase different from that of A_0 we obtain:

$$\begin{aligned}
M_{\bar{B}_d \rightarrow \pi^+ \pi^-} &= \frac{G_F}{\sqrt{2}} |V_{ub} V_{ud}^*| m_B^2 f_\pi f_+(0) \left\{ e^{-i\gamma} \frac{1}{2\sqrt{3}} A_2 e^{i\delta_2} + \right. \\
&\quad \left. + e^{-i\gamma} \frac{1}{\sqrt{6}} A_0 e^{i\delta_0} + \left| \frac{V_{td}^* V_{tb}}{V_{ub} V_{ud}^*} \right| e^{i\beta} P e^{i(\delta_p + \tilde{\delta}_0)} \right\} , \quad (1)
\end{aligned}$$

$$\begin{aligned}
M_{\bar{B}_d \rightarrow \pi^0 \pi^0} &= \frac{G_F}{\sqrt{2}} |V_{ub} V_{ud}^*| m_B^2 f_\pi f_+(0) \left\{ e^{-i\gamma} \frac{1}{\sqrt{3}} A_2 e^{i\delta_2} - \right. \\
&\quad \left. - e^{-i\gamma} \frac{1}{\sqrt{6}} A_0 e^{i\delta_0} - \left| \frac{V_{td}^* V_{tb}}{V_{ub} V_{ud}^*} \right| e^{i\beta} P e^{i(\delta_p + \tilde{\delta}_0)} \right\} , \quad (2)
\end{aligned}$$

$$M_{\bar{B}_u \rightarrow \pi^- \pi^0} = \frac{G_F}{\sqrt{2}} |V_{ub} V_{ud}^*| m_B^2 f_\pi f_+(0) \left\{ \frac{\sqrt{3}}{2\sqrt{2}} e^{-i\gamma} A_2 e^{i\delta_2} \right\} , \quad (3)$$

where V_{ik} are CKM matrix elements and the penguin amplitude with an intermediate c -quark multiplied by $V_{ub} V_{ud}^* + V_{cb} V_{cd}^* + V_{tb} V_{td}^* = 0$ is subtracted from the penguin amplitudes with intermediate u -, c - and t -quarks (the so-called t -convention)². To check if the factorization works in $B \rightarrow \pi\pi$ decays it is convenient to introduce $f_+(0)$ - the value of the formfactor which enters the amplitude of semileptonic $B_d \rightarrow \pi l \nu$ decay at zero momentum transfer in Eqs. (1)-(3). γ and β are the angles of the unitarity triangle; δ_2 and δ_0 are FSI phases of the tree amplitudes with $I = 2$ and $I = 0$ (below we will use $\delta \equiv \delta_2 - \delta_0$), δ_p originates from the imaginary part of the penguin loop with c -quark propagating in it [5] while $\tilde{\delta}_0$ is long distance FSI phase of the penguin amplitude. $\tilde{\delta}_0$ in general is different from δ_0 ; in Section 4 we will argue that $\rho\rho$ intermediate state generate large value of δ_0 while its contribution into $\tilde{\delta}_0$ is smaller: (pseudo)scalar part of penguin operator do not produce ρ mesons.

The charge conjugate amplitudes are obtained by the same formulas with substitution $\beta, \gamma \rightarrow -\beta, -\gamma$.

The CP asymmetries are given by [6]:

$$C_{\pi\pi} \equiv \frac{1 - |\lambda_{\pi\pi}|^2}{1 + |\lambda_{\pi\pi}|^2} , \quad S_{\pi\pi} \equiv \frac{2\text{Im}(\lambda_{\pi\pi})}{1 + |\lambda_{\pi\pi}|^2} , \quad \lambda_{\pi\pi} \equiv e^{-2i\beta} \frac{M_{\bar{B} \rightarrow \pi\pi}}{M_{B \rightarrow \pi\pi}} ,$$

where $\pi\pi$ is $\pi^+ \pi^-$ or $\pi^0 \pi^0$.

²We prefer t -convention since the penguin contribution to the amplitude A_0 in it is suppressed as $m_c^2/(m_b^2 \ln m_W^2/m_b^2)$, while in the c -convention one subtracts the penguin amplitude with intermediate t -quark thus making the penguin contribution to A_0 comparable to the penguin term.

Now we have all the necessary formulas and neglecting the penguin contribution we are able to determine A_0 , A_2 , δ and the value of the unitarity triangle angle α from the experimental data on B_{+-} , B_{00} , B_{+0} and S_{+-} , which are presented in Table 1. By definition:

$$\begin{aligned} B_{+-} &\equiv 1/2[\text{Br}(B_d \rightarrow \pi^+\pi^-) + \text{Br}(\bar{B}_d \rightarrow \pi^+\pi^-)] , \\ B_{00} &\equiv 1/2[\text{Br}(B_d \rightarrow \pi^0\pi^0) + \text{Br}(\bar{B}_d \rightarrow \pi^0\pi^0)] , \\ B_{+0} &= \text{Br}(B_u \rightarrow \pi^+\pi^0) = \text{Br}(\bar{B}_u \rightarrow \pi^-\pi^0) , \end{aligned}$$

the last equality holds as far as the electroweak penguins are neglected.

To extract the product $A_2 f_+(0)$ from B_{+0} we will use the value of $|V_{ub}|$ obtained from the general fit of the Wolfenstein parameters of CKM matrix (CKM fitter, summer 2005): $A = 0.825 \pm 0.019$, $\lambda = 0.226 \pm 0.001$, $\bar{\rho} = 0.207 \pm 0.040$, $\bar{\eta} = 0.340 \pm 0.023$:

$$|V_{ub}| = (3.90 \pm 0.10) \cdot 10^{-3} . \quad (4)$$

From (3) and the experimental data on B_{+0} from the last column of Table 1 we readily get:

$$A_2 f_+(0) = 0.35 \pm 0.02 . \quad (5)$$

In order to understand if the factorization works in $B_u \rightarrow \pi^+\pi^0$ decay we should determine the value of $f_+(0)$. We find it using the data on $B \rightarrow \pi l \nu$ decay from [8]:

$$f_+(0) = 0.22 \pm 0.02 , \quad (6)$$

thus getting:

$$A_2 = 1.60 \pm 0.20 , \quad (7)$$

which is not far from the result of factorization:

$$A_2^f = \frac{8}{3\sqrt{3}}(c_1 + c_2) \equiv \frac{2}{\sqrt{3}}(a_1 + a_2) = 1.35 , \quad (8)$$

see Appendix. We come to the same conclusion as the authors of paper [9]: A_2 is estimated correctly by factorization. Neglecting the penguin contribution we are able to extract the values of A_0 and FSI phases difference δ from Eqs. (1)-(3) and the experimental data for B_{+-} , B_{00} and B_{+0} from the last column of Table 1. In this way we obtain:

$$A_0 = 1.53 \pm 0.23 , \quad (9)$$

which should be compared with the result of factorization:

$$A_0^f = \frac{\sqrt{2}}{3\sqrt{3}}(5c_1 - c_2) = 1.54 \quad , \quad (10)$$

see Appendix. In this way we come to the conclusion that factorization works well for the moduli of both decay amplitudes.

For the phase difference $\delta \equiv \delta_2 - \delta_0$ we get:

$$\cos \delta = \frac{\sqrt{3}}{4} \frac{B_{+-} - 2B_{00} + \frac{2}{3} \frac{\tau_0}{\tau_+} B_{+0}}{\sqrt{\frac{\tau_0}{\tau_+} B_{+0}} \sqrt{B_{+-} + B_{00} - \frac{2}{3} \frac{\tau_0}{\tau_+} B_{+0}}} \quad , \quad (11)$$

$$\delta = \pm(53^\circ \pm 7^\circ) \quad , \quad (12)$$

where $\tau_0/\tau_+ \equiv \tau(B_d)/\tau(B_u) = 0.92$ is substituted. This is the place where the factorization which predicts the negligible FSI phases fails.

In Section 4 we will present a model in which the pattern of $B \rightarrow \pi\pi$ amplitudes obtained above is realized.

Let us turn to the bottom part of Table 1. Since we neglect penguins the experimental value of S_{+-} is directly related to the unitarity triangle angle α :

$$\sin 2\alpha^T = S_{+-} \quad , \quad (13)$$

$$\alpha_{\text{BABAR}}^T = 99^\circ \pm 5^\circ \quad , \quad \alpha_{\text{Belle}}^T = 111^\circ \pm 6^\circ \quad , \quad \alpha_{\text{average}}^T = 105^\circ \pm 4^\circ \quad ,$$

where index “T” stands for “tree” stressing that penguins are neglected (three other values of α are not compatible with the Standard Model).

3 FSI phases in $K \rightarrow \pi\pi$, $D \rightarrow \pi\pi$ and $B \rightarrow D\pi$

The s -wave amplitudes of two pions production with $I = 0$ and $I = 2$ are generally different. In particular there are quark-antiquark resonances in s -channel with $I = 0$ but not with $I = 2$. This can lead to large difference of phases in the channels with $I = 0$ and $I = 2$. Let us remind what experimental data tell us about these phases at the pion center of mass energies $E = m_K$ and $E = m_D$. Since at $E = m_K$ only elastic rescattering of pions is possible (the inelastic channels are closed since the energy is low) Watson theorem is applicable and strong interaction phases of matrix elements of $K \rightarrow (\pi\pi)_I$ decays are equal to the phases of amplitudes describing $\pi\pi \rightarrow \pi\pi$ scattering at

$E = m_K$. From the analysis of $\pi\pi \rightarrow \pi\pi$ scattering data performed in [10] at $E_{\pi\pi} = m_K$ we have:

$$\delta_0^K = 35^\circ \pm 3^\circ , \quad (14)$$

$$\delta_2^K = -7^\circ \pm 0.2^\circ , \quad (15)$$

$$\delta_0^K - \delta_2^K = 42^\circ \pm 4^\circ . \quad (16)$$

The large value of δ_0^K is due to the specific behaviour of $I = 0, J = 0$ $\pi\pi$ -phase attributed to the $f_0(600)$ (or σ) “resonance”.

The same value of the difference $\delta_0^K - \delta_2^K$ follows from the analysis of $K_S \rightarrow \pi^+\pi^-$, $K_S \rightarrow \pi^0\pi^0$ and $K^+ \rightarrow \pi^+\pi^0$ decay probabilities analogous to one we perform for $B \rightarrow \pi\pi$ decays in Section 2 neglecting the penguin contributions. (In case of $K \rightarrow \pi\pi$ decays the penguins are very important being responsible for the enhancement of $I = 0$ amplitude. Since CKM phase of the penguin amplitudes is almost the same as that of the tree amplitude, the analysis performed in Section 2 is applicable for kaon decays but the amplitude A_0^K should contain the penguin contribution as well.)

What concerns the moduli of the kaon decay amplitudes with $I = 0$ and $I = 2$, they are given with rather good accuracy (within 50% from the experimental data) by factorization [11].

In case of $D \rightarrow \pi\pi$ decays, the gluon penguin amplitudes are negligible in comparison with the tree ones (since the loop with s -quark is subtracted from the one with d -quark while the momentum transfer is of the order of m_D^2), and the effective Hamiltonian responsible for these decays looks like:

$$\begin{aligned} \hat{H}_D = & \frac{G_F}{\sqrt{2}} \sin \theta_c [c_1^D \bar{u} \gamma_\alpha (1 + \gamma_5) d d \bar{d} \gamma_\alpha (1 + \gamma_5) c + \\ & + c_2^D \bar{d} \gamma_\alpha (1 + \gamma_5) d \bar{u} \gamma_\alpha (1 + \gamma_5) c] , \end{aligned} \quad (17)$$

where θ_c is Cabibbo angle, $\sin \theta_c = 0.22$, $c_1^D \approx 1.22$, $c_2^D \approx -0.42$ [12].

Calculating the matrix elements in the factorization approximation we obtain:

$$M_{D \rightarrow \pi^+\pi^-} = \frac{G_F}{\sqrt{2}} \sin \theta_c f_+^D(0) f_\pi m_D^2 (c_1^D + \frac{c_2^D}{3}) , \quad (18)$$

$$M_{D \rightarrow \pi^0\pi^0} = \frac{G_F}{\sqrt{2}} \sin \theta_c f_+^D(0) f_\pi m_D^2 (\frac{c_1^D}{3} + c_2^D) , \quad (19)$$

$$M_{D^\pm \rightarrow \pi^\pm\pi^0} = \frac{G_F}{\sqrt{2}} \sin \theta_c f_+^D(0) \frac{f_\pi}{\sqrt{2}} m_D^2 \frac{4}{3} (c_1^D + c_2^D) , \quad (20)$$

(the analogous formulas for B decays are derived in Appendix) while the isotopic analysis gives:

$$M_{D \rightarrow \pi^+ \pi^-} = \frac{G_F}{\sqrt{2}} \sin \theta_c f_+^D(0) f_\pi m_D^2 \left(\frac{e^{i\delta_d}}{2\sqrt{3}} A_2^D + \frac{1}{\sqrt{6}} A_0^D \right) , \quad (21)$$

$$M_{D \rightarrow \pi^0 \pi^0} = \frac{G_F}{\sqrt{2}} \sin \theta_c f_+^D(0) f_\pi m_D^2 \left(\frac{e^{i\delta_d}}{\sqrt{3}} A_2^D - \frac{1}{\sqrt{6}} A_0^D \right) , \quad (22)$$

$$M_{D^\pm \rightarrow \pi^\pm \pi^0} = \frac{G_F}{\sqrt{2}} \sin \theta_c f_+^D(0) f_\pi m_D^2 \left(\frac{e^{i\delta_d} \sqrt{3}}{2\sqrt{2}} A_2^D \right) . \quad (23)$$

From the recent study of semileptonic D -meson decays $D \rightarrow \pi l \nu$ it was found [13]:

$$f_+^D(0) = 0.62 \pm 0.04 . \quad (24)$$

Comparing (23), (24) and recent measurement [14]:

$$\text{Br}(D^\pm \rightarrow \pi^\pm \pi^0) = (1.25 \pm 0.10) \cdot 10^{-3} \quad (25)$$

we obtain:

$$A_2^D = 0.88 \pm 0.08 ,$$

which is not so different from the factorization result, (20):

$$(A_2^D)_f = \frac{8}{3\sqrt{3}} (c_1^D + c_2^D) = 1.2 . \quad (26)$$

Comparing Eqs. (18)-(19) with Eqs. (21)-(22) we obtain in the factorization approximation:

$$(A_0^D)_f = \frac{\sqrt{2}}{3\sqrt{3}} (5c_1^D - c_2^D) = 1.8 , \quad (27)$$

while according to [14] from the experimental data it follows:

$$A_0^D = 2.1 \pm 0.2 . \quad (28)$$

We see that the factorization results are within 30% from the experimental values of the moduli of the decay amplitudes. However, factorization fails completely in describing the difference of FSI phases. The data on $D \rightarrow \pi^+ \pi^-$, $D \rightarrow \pi^0 \pi^0$ and $D^\pm \rightarrow \pi^\pm \pi^0$ branching ratios lead to [14]:

$$\delta_2^D - \delta_0^D \equiv \delta_D = \pm(86^\circ \pm 4^\circ) , \quad (29)$$

which is responsible for (or follows from) the relatively large $D \rightarrow \pi^0 \pi^0$ decay probability [14]:

$$\text{Br}(D^0 \rightarrow \pi^0 \pi^0)_{\text{exp}} = (0.79 \pm 0.08) \cdot 10^{-3} , \quad (30)$$

of the order of that into $\pi^+\pi^-$ [15]:

$$\text{Br}(D^0 \rightarrow \pi^+\pi^-)_{\text{exp}} = (1.39 \pm 0.07) \cdot 10^{-3} .$$

Using $\tau_{D_0}/\tau_{D_+} = 410/1040$ from [15] we readily reproduce the phase difference given by (29) with the help of (11).

In factorization approximation neglecting δ_D we will get:

$$\begin{aligned} \text{Br}(D^0 \rightarrow \pi^0\pi^0)_f &= \frac{1}{2} \left[\frac{\frac{1}{3}c_1^D + c_2^D}{\frac{2\sqrt{2}}{3}(c_1^D + c_2^D)} \right]^2 \times \\ &\times \text{Br}(D^+ \rightarrow \pi^\pm\pi^0) \frac{\tau_{D_0}}{\tau_{D_+}} \ll \text{Br}(D^0 \rightarrow \pi^0\pi^0)_{\text{exp}} . \end{aligned} \quad (31)$$

The analogous phenomena we encountered in $B \rightarrow \pi\pi$ decays.

Let us note that the s -wave resonance with zero isospin $f_0(1710)$ alone cannot explain such a big phase; its contribution to $D \rightarrow (\pi\pi)_{I=0}$ decay amplitude is proportional to:

$$m_D - m_{f_0} - i\frac{\Gamma_{f_0}}{2} = 150 \text{ MeV} - i70 \text{ MeV} , \quad |\delta_D| \approx 30^\circ . \quad (32)$$

It is not easy to reconcile reasonable (20%÷30%) accuracy of factorization in describing the moduli of the decay amplitudes into $\pi\pi$ states with a definite isospin and the large FSI phases difference since the latter signal of strong rescattering of pions at $E \approx m_D$ which should not only generate phases but also shift the moduli of the amplitudes. The resolution may be that the interactions are “semistrong” in both channels: one half of 86° comes from $I = 0$, another from $I = 2$ (just as in the case of $B \rightarrow \pi\pi$ decays, see Section 4).

If we suppose that FSI phases scale with decaying meson mass as $1/M$ we will get about 30° phases difference for $B \rightarrow \pi\pi$ decays from (29).

$\pi\pi$ FSI phase shifts at $E = m_K$ and $E = m_D$ are not small. However, in both cases we are in the regions where two pion resonances are situated, which is not the case for the high energy of the order of B -meson mass.

Our last example is $B \rightarrow D\pi$ decays, where the energy is high and we are definitely above the resonances domain, though the FSI phase shift is nevertheless large³. $D\pi$ pair produced in B -decays can have $I = 1/2$ or $3/2$. From the measurement of the probabilities of

³We are grateful to A.E. Bondar who brought this case to our attention.

$B^- \rightarrow D^0\pi^-$, $B^0 \rightarrow D^-\pi^+$ and $B^0 \rightarrow D^0\pi^0$ decays in paper [16] the FSI phases difference of these two amplitudes was determined:

$$\delta_{D\pi} = 30^\circ \pm 7^\circ . \quad (33)$$

Concluding this section we wish to note that the direct CP asymmetry observed in $B_d(\bar{B}_d) \rightarrow \pi^\mp K^\pm$ decays is incompatible with small FSI phase difference between $I = 1/2$ and $I = 3/2$ amplitudes.

4 FSI phases: theoretical considerations

There are many theoretical papers on the final state interaction (FSI) in the heavy-meson decays [17] - [22]. For example in paper [18] the final state interactions in $B \rightarrow \pi\pi$ decay are modelled as the soft rescattering of the certain intermediate two-body hadronic channels ($\pi\pi, \rho\rho, D^*\bar{D}^*, D\bar{D}$). The hadronic amplitudes, which enter the calculation of the imaginary parts of the decay amplitudes were described by π, ρ, D, D^* -meson exchanges in the t -channel. Rather large phases due to FSI have been obtained. While for π -exchange (with the pole close to the physical region) this procedure is reasonable, it exaggerates the contributions of the vector exchanges (ρ, D^*), which for the elementary particle exchange with spin $J = 1$ gives the partial wave amplitude, which does not decrease with energy. In reality all the exchanges should be reggeized and in the physical region of the processes corresponding intercepts $\alpha_i(0) < 1$ (for D^* the most probably value of the intercept is negative). This will lead to the strong reduction of the corresponding amplitudes (see for example [19]).

A number of papers ([19] - [22]) use Watson theorem in order to extract the phases of the decay amplitudes by multiplying the bare matrix elements by $S^{1/2}$ (where S is the S -matrix). However in B -decays there are many coupled multiparticle channels. In this case such a procedure can be applied only in the basis of the eigenstates which diagonalize S -matrix. But for the realistic strong interactions this is impossible at present.

Another approximation is to use the Feynman diagrams approach taking only the low mass intermediate states X, Y into account. This approach coincides with the use of the unitarity condition only if the transitions $\pi\pi \rightarrow XY$ are described by the real amplitudes. This is certainly not true for elastic $\pi\pi$ -scattering, where the amplitude

at large energies is predominantly imaginary. In this formalism the resulting decay matrix elements are:

$$M_{\pi\pi}^I = M_{XY}^{(0)I} (\delta_{\pi X} \delta_{\pi Y} + iT_{XY \rightarrow \pi\pi}^{J=0}) \quad , \quad (34)$$

where $M_{XY}^{(0)I}$ are the decay matrix elements without FSI and $T_{XY \rightarrow \pi\pi}^{J=0}$ is the $J = 0$ partial wave amplitude of the process $XY \rightarrow \pi\pi$ ⁴. At very high energies the amplitudes of $\pi\pi$ elastic scattering are imaginary and $T_{XY \rightarrow \pi\pi}^{J=0}$ do not decrease with energy (mass of a heavy meson). Thus this contribution, according to (34) does not change the phase of the matrix element, but only changes its modulus. The extra phases come from the real parts of the amplitudes, which in Regge model are due to the secondary exchanges ($R \equiv \rho, f, \dots$), which decrease with energy as $1/s^{1-\alpha_R(0)} \approx 1/\sqrt{s}$. The contribution of the pion exchange in the t -channel, which is dominant in the process $\rho\rho \rightarrow \pi\pi$ ⁵ decreases even faster (as $1/s$). However $\text{Br}(B \rightarrow \rho^+\rho^-)$ is substantially larger than $\text{Br}(B \rightarrow \pi^+\pi^-)$ and $\rho\rho$ intermediate state is important in (34). This is especially true for color suppressed $B \rightarrow \pi^0\pi^0$ decay, where the chain $B \rightarrow \rho^+\rho^- \rightarrow \pi^0\pi^0$ is enhanced. On the contrary $\pi\pi$ contribution in $B \rightarrow \rho\rho$ decay is relatively suppressed. Using Regge analysis of $\pi\pi$ scattering [23] and π -exchange model for $\rho\rho \rightarrow \pi\pi$ transitions, we obtain phases due to final state interactions for $\pi\pi$ final state $\delta_2 \approx -12^\circ$ and $\delta_0 \approx 18^\circ$. Thus the phase difference $\delta \approx -30^\circ$ is generated by intermediate $\rho\rho$ and $\pi\pi$ states⁶.

Sign of δ is negative, just as in the case of $K \rightarrow \pi\pi$ decays. In this way in the numerical estimates we will use negative value of δ from (12):

$$\delta \approx -(50 \pm 7)^\circ \quad , \quad \delta_0 = 30^\circ \quad , \quad \delta_2 = -20^\circ \quad . \quad (35)$$

As far as $\text{Br}(B \rightarrow \rho^+\rho^-)$ is much larger than $\text{Br}(B \rightarrow \pi^+\pi^-)$ because of enhancement in tree amplitudes but not in penguin amplitudes (contribution of penguins in $B \rightarrow \rho\rho$ amplitude is small) we should expect $\tilde{\delta}_0$ to be substantially smaller than δ_0 .

Note that in this model there is little change in moduli of amplitudes in comparison with factorization predictions.

For $B \rightarrow \rho\rho$ decays the same model gives $\approx -5^\circ$ for $I = 2$ and $+5^\circ$ for $I = 0$ amplitudes, resulting in a small phase difference $\approx 10^\circ$,

⁴We use the standard normalization with $T^J = \frac{S^J - 1}{2i}$.

⁵In B -decays transverse polarizations of ρ -mesons are small that is why a_2 and ω exchanges in $\rho\rho \rightarrow \pi\pi$ amplitudes are suppressed.

⁶An accuracy of this number is about 15° .

consistent with experiment.

Thus the lowest mass hadronic intermediate states may produce the phases which are consistent with the data on $B \rightarrow \pi\pi$ and $B \rightarrow \rho\rho$ decays. There are many high-mass states as well, and they can lead to additional phases (the inclusion of πa_1 intermediate state makes $\delta \approx -40^\circ$).

5 Taking penguins into account: shifts of A_0 , δ and α and the values of C_{+-} and C_{00}

Let us analyse to what changes of the parameters introduced and calculated in Section 2 penguins lead. Since QCD penguins contribute only to $I = 0$ amplitude the value of A_2 extracted from B_{+0} remains the same, see (7). The requirement that the numerical values of B_{+-} and B_{00} are not shifted when penguins are taken into account leads to the following shifts of the amplitude A_0 and phase difference δ :

$$A_0 \rightarrow A_0 + \tilde{A}_0, \quad \delta \rightarrow \delta + \tilde{\delta}, \quad (36)$$

$$\tilde{A}_0 = \sqrt{6} \left| \frac{V_{td}}{V_{ub}} \right| \cos \alpha \cos(\delta_p + \tilde{\delta}_0 - \delta_0) P, \quad (37)$$

$$\tilde{\delta} = -\sqrt{6} \left| \frac{V_{td}}{V_{ub}} \right| \cos \alpha \sin(\delta_p + \tilde{\delta}_0 - \delta_0) P / A_0, \quad (38)$$

where only the terms linear in P are taken into account. For numerical estimates we take:

$$\left| \frac{V_{td}}{V_{ub}} \right| = \frac{\sin \gamma}{\sin \beta} \approx 2.3 \pm 0.2, \quad (39)$$

where $\beta = 22^\circ$, $\gamma = 60^\circ \pm 10^\circ$. In the factorization approach we have (see Appendix)⁷:

$$P^f = -a_4 - \frac{2m_\pi^2}{(m_u + m_d)m_b} a_6 = 0.06, \quad (40)$$

and shifts of A_0 and δ are small:

$$-0.12 < \tilde{A}_0 < 0.12, \quad -4^\circ < \tilde{\delta} < 4^\circ \quad (41)$$

for

$$A_0 = 1.5, \quad -1 < \cos(\delta_p + \tilde{\delta}_0 - \delta_0), \quad \sin(\delta_p + \tilde{\delta}_0 - \delta_0) < 1 \quad \text{and} \quad 70^\circ < \alpha < 110^\circ. \quad (42)$$

⁷Note that the definition of P used in the present paper differs in sign from that in [4].

In particular even if the penguin contribution is underestimated by factor 2, the statement that $\delta + \tilde{\delta}$ is large still holds⁸ (note that α can be closer to 90°).

The following two equations for direct CP asymmetries determine P and $\delta_p + \tilde{\delta}_0 - \delta_0$ (as far as A_0 , A_2 and $\delta = \delta_2 - \delta_0$ are known):

$$C_{+-} = -\frac{\tilde{P}}{\sqrt{3}} \sin \alpha [\sqrt{2} A_0 \sin(\delta_0 - \tilde{\delta}_0 - \delta_p) + A_2 \sin(\delta_2 - \tilde{\delta}_0 - \delta_p)] /$$

$$/ \left[\frac{A_0^2}{6} + \frac{A_2^2}{12} + \frac{A_0 A_2}{3\sqrt{2}} \cos \delta - \sqrt{\frac{2}{3}} A_0 \tilde{P} \cos \alpha \cos(\delta_0 - \tilde{\delta}_0 - \delta_p) - \right.$$

$$\left. - \frac{A_2 \tilde{P}}{\sqrt{3}} \cos \alpha \cos(\delta_2 - \tilde{\delta}_0 - \delta_p) + \tilde{P}^2 \right] , \quad (43)$$

$$C_{00} = -\sqrt{\frac{2}{3}} \tilde{P} \sin \alpha [A_0 \sin(\delta_0 - \tilde{\delta}_0 - \delta_p) - \sqrt{2} A_2 \sin(\delta_2 - \tilde{\delta}_0 - \delta_p)] /$$

$$/ \left[\frac{A_0^2}{6} + \frac{A_2^2}{3} - \frac{\sqrt{2}}{3} A_0 A_2 \cos \delta - \sqrt{\frac{2}{3}} A_0 \tilde{P} \cos \alpha \cos(\delta_0 - \tilde{\delta}_0 - \delta_p) + \right.$$

$$\left. + \frac{2}{\sqrt{3}} A_2 \tilde{P} \cos \alpha \cos(\delta_2 - \tilde{\delta}_0 - \delta_p) + \tilde{P}^2 \right] , \quad (44)$$

where

$$\tilde{P} \equiv \left| \frac{V_{td}^* V_{tb}}{V_{ub} V_{ud}^*} \right| P \approx 2.3 P . \quad (45)$$

Three last terms in denominators of (43) and (44) lead to less than 10% variations of the numerical values of C_{+-} and C_{00} for $\tilde{P} < 0.3$. Neglecting them we get:

$$\frac{\sin(\delta_2 - \tilde{\delta}_0 - \delta_p)}{\sin(\delta_0 - \tilde{\delta}_0 - \delta_p)} = \frac{1 - 0.57 \frac{C_{00}}{C_{+-}}}{1.4 + 0.41 \frac{C_{00}}{C_{+-}}} , \quad (46)$$

where the numerical values for A_2 , A_0 and δ from (7), (9) and (12) correspondingly were used.

From the central values in the last column in Table 1 of C_{+-} and C_{00} we get:

$$\delta_0 - \tilde{\delta}_0 - \delta_p = 70^\circ , \quad P = 0.11 . \quad (47)$$

The numerical value of P is two times larger than the factorization estimate of it presented in (40), while $\delta_0 - \tilde{\delta}_0 - \delta_p$ largely deviates from

⁸Indication of such an underestimate follows from the probability of $b \rightarrow s$ penguin dominated $B^+ \rightarrow K^0 \pi^+$ decay.

30° which is our estimate of δ_0 , while $\tilde{\delta}_0$ should be considerably smaller as well as δ_p the latter being close to 30° only for very asymmetric configurations of quarks in π mesons and is smaller otherwise [5]. If the experimental accuracy of C_{ik} were good we would be able to use the results obtained for determination of the value of the angle α from S_{+-} , realizing in this way Gronau-London approach [1].

However the experimental uncertainty in C_{00} is very big, while the measurements of C_{+-} by Belle and BABAR contradict each other. So let us look which values of the direct asymmetries follow from our formulas.

Denominator of the expression for C_{+-} is close to one and in the expression in brackets in nominator first term dominates. Neglecting $\tilde{\delta}_0$ and δ_p and taking $\delta_0 = 30^\circ$, $\delta_2 = -20^\circ$ we get $C_{+-} = -0.04$ for the value of penguin amplitude obtained in the factorization approach, (40). We reproduce BABAR central value of C_{+-} if we suppose that factorization underestimate penguin amplitude by factor 2; however in order to reproduce Belle number we should accept that factorization is wrong by factor 10, which looks highly improbable.

What to do if C_{+-} appeared to be equal to the average of the present day Belle and BABAR results $C_{+-} \approx -0.3$? One possibility is to suppose that $\delta_2 - \tilde{\delta}_0 - \delta_P \approx 0$, while $\delta_0 - \tilde{\delta}_0 - \delta_P \approx 50^\circ$ and to look for FSI mechanism which provides such a result⁹.

C_{00} is also negative while its absolute value is larger than C_{+-} : denominator is about .55 while in nominator both terms are negative.

The requirement that the value of CP asymmetry S_{+-} is not changed when penguins are taken into account leads to the following shift of the value of the unitarity triangle angle α :

$$\alpha = \alpha^T + \tilde{\alpha} \quad , \quad (48)$$

$$\tilde{\alpha} = -\frac{\tilde{P}}{2\sqrt{3}} \sin \alpha [\sqrt{2}A_0 \cos(\delta_0 - \tilde{\delta}_0 - \delta_P) + A_2 \cos(\delta_2 - \tilde{\delta}_0 - \delta_P)] \quad . \quad (49)$$

Substituting the numerical values of A_2 from (7), A_0 from (9), $\tilde{P}$ from (45) and substituting $\sin \alpha$ by one and both $\cos$ by 0.9 we get:

$$\tilde{\alpha} \approx -2.3P \approx -7^\circ, \quad (50)$$

where the result of the matrix element of the penguin operator calculation in factorization approximation (40) is used. We observe that

⁹Let us note that in paper [24] argument in favor of $C_{+-} \approx -0.3$ is presented which is based on the comparison of the direct CP violation in $B(\bar{B}) \rightarrow \pi^+\pi^-$ and $B(\bar{B}) \rightarrow K^+\pi^-(K^-\pi^+)$ decays (see also [25]).

our approach is at least selfconsistent: the shift of α due to penguin contribution is small. For the BABAR value of S_{+-} we obtain:

$$\alpha_{\text{BABAR}} = \alpha_{\text{BABAR}}^T + \tilde{\alpha} = 92^\circ \pm 5^\circ . \quad (51)$$

In the case of the averaged experimental values we get:

$$\alpha_{\text{average}} = \alpha_{\text{average}}^T + \tilde{\alpha} = 98^\circ \pm 4^\circ . \quad (52)$$

Theoretical uncertainty of the value of α can be estimated in the following way. Let us suppose that the accuracy of the factorization calculation of the penguin amplitude is 100% (in all the examples considered in this paper it was much better). Then:

$$\tilde{\alpha} = -7^\circ \pm 7_{\text{theor}}^\circ , \quad (53)$$

$$\alpha_{\text{average}} = 98^\circ \pm 4_{\text{exp}}^\circ \pm 7_{\text{theor}}^\circ , \quad (54)$$

while BABAR value is smaller:

$$\alpha_{\text{BABAR}} = 92^\circ \pm 5_{\text{exp}}^\circ \pm 7_{\text{theor}}^\circ . \quad (55)$$

Better theoretical accuracy of α follows from $B \rightarrow \rho^+ \rho^-$ decays, where penguin contribution is two times smaller. Since FSI phases are small in these decays, results of the paper [4] are directly applicable:

$$\alpha^{\rho\rho} = 92^\circ \pm 7_{\text{exp}}^\circ \pm 4_{\text{theor}}^\circ , \quad (56)$$

where we take the WHOLE penguin contribution as an estimate of the theoretical uncertainty.

The model independent isospin analysis of $B \rightarrow \rho\rho$ decays performed by BABAR gives [26] :

$$\alpha_{\text{BABAR}}^{\rho\rho} = 100^\circ \pm 13^\circ , \quad (57)$$

while the analogous analysis performed by Belle gives [27] :

$$\alpha_{\text{Belle}}^{\rho\rho} = 87^\circ \pm 17^\circ . \quad (58)$$

Finally, the global CKM fit results are[28, 29]:

$$\alpha_{\text{CKMfitter}} = 97^\circ \pm 5^\circ , \quad \alpha_{\text{UTfit}} = 95^\circ \pm 5^\circ . \quad (59)$$

6 Conclusions

1. The moduli of the amplitudes A_0 and A_2 of B decays into $\pi\pi$ states are given with good accuracy by factorization of the tree quark diagram, while FSI phase shift between these two amplitudes is very large, $|\delta| \approx 50^\circ$, which explains large $B_d \rightarrow \pi^0\pi^0$ decay probability.
2. Theoretical uncertainty of the value of α extracted from $B \rightarrow \pi\pi$ data on S_{+-} is at the level of few degrees.
3. Resolution of the contradiction of Belle and BABAR experimental data on CP asymmetries in $B \rightarrow \pi\pi$ decays is very important both for understanding the FSI dynamics (the data on C) and for determination of angle α (the data on S).

We are grateful to A.E. Bondar, A.Yu. Khodjamirian and L.B. Okun for useful discussions.

A.K. was partly supported by grants CRDF RUP 2-2621-M0-04 and RFBR 04-02-17263; M.V. was partly supported by grants RFBR 05-02-17203 and NSh-5603.2006.2.

Table 1. Experimental data on $B \rightarrow \pi\pi$ decays. Branching ratios are in units of 10^{-6} .

	BABAR	Belle	Heavy Flavor Averaging Group [7]
B_{+-}	5.5 ± 0.5	4.4 ± 0.7	5.0 ± 0.4
B_{00}	1.17 ± 0.33	2.3 ± 0.5	1.45 ± 0.29
B_{+0}	5.8 ± 0.7	5.0 ± 1.3	5.5 ± 0.6
S_{+-}	-0.30 ± 0.17	-0.67 ± 0.16	-0.50 ± 0.12
C_{+-}	-0.09 ± 0.15	-0.56 ± 0.13	-0.37 ± 0.10
C_{00}	-0.12 ± 0.56	-0.44 ± 0.56	-0.28 ± 0.39

Appendix

In order to calculate amplitude A_2 in the factorization approximation we start from the following Hamiltonian:

$$\begin{aligned}\hat{H} = & \frac{G_F}{\sqrt{2}} |V_{ub}V_{ud}^*| e^{-i\gamma} [c_1 \bar{u}\gamma_\alpha(1 + \gamma_5)b\bar{d}\gamma_\alpha(1 + \gamma_5)u + \\ & + c_2 \bar{d}\gamma_\alpha(1 + \gamma_5)b\bar{u}\gamma_\alpha(1 + \gamma_5)u] ,\end{aligned}\quad (\text{A1})$$

and take the matrix element of it between $\bar{B}_u$ and $\pi^-\pi^0$ states:

$$\begin{aligned}M_{\bar{B}_u \rightarrow \pi^-\pi^0}^f &= \frac{G_F}{\sqrt{2}} |V_{ub}V_{ud}^*| e^{-i\gamma} \left[(c_1 + \frac{c_2}{3}) \langle \pi^- | \bar{d}\gamma_\alpha(1 + \gamma_5)u | 0 \rangle \times \right. \\ &\times \langle \pi^0 | \bar{u}\gamma_\alpha(1 + \gamma_5)b | \bar{B}_u \rangle + (\frac{1}{3}c_1 + c_2) \langle \pi^0 | \bar{u}\gamma_\alpha(1 + \gamma_5)u | 0 \rangle \times \\ &\times \left. \langle \pi^- | \bar{d}\gamma_\alpha(1 + \gamma_5)b | \bar{B}_u \rangle \right] = \\ &= \frac{G_F}{\sqrt{2}} |V_{ub}V_{ud}^*| e^{-i\gamma} \frac{4}{3} (c_1 + c_2) \langle \pi^- | \bar{d}\gamma_\alpha(1 + \gamma_5)u | 0 \rangle \times \\ &\times \langle \pi^0 | \bar{u}\gamma_\alpha(1 + \gamma_5)b | \bar{B}_u \rangle = \frac{G_F}{\sqrt{2}} |V_{ub}V_{ud}^*| e^{-i\gamma} \frac{4}{3} (c_1 + c_2) f_\pi P_\alpha^{\pi^-} \times \\ &\times \frac{1}{\sqrt{2}} f_+(0) (P_\alpha^B + P_\alpha^{\pi^0}) = \\ &= \frac{G_F}{\sqrt{2}} |V_{ub}V_{ud}^*| e^{-i\gamma} f_\pi m_B^2 \frac{1}{\sqrt{2}} f_+(0) \frac{4}{3} (c_1 + c_2) .\end{aligned}\quad (\text{A2})$$

Comparing the last expression with (3) we get:

$$A_2^f = \frac{8}{3\sqrt{3}} (c_1 + c_2) = \frac{2}{\sqrt{3}} (a_1 + a_2) = 1.35 , \quad (\text{A3})$$

where $a_1 \equiv c_1 + c_2/3 = 1.02$, $a_2 \equiv c_2 + c_1/3 = 0.15$ and the values $c_1 = 1.09$, $c_2 = -0.21$ from paper [12] were used.

To find the amplitude A_0 in the factorization approximation let us take the matrix element of $\hat{H}$ between $\bar{B}_d$ and $\pi^+\pi^-$ -states:

$$M_{\bar{B}_d \rightarrow \pi^+\pi^-}^f = \frac{G_F}{\sqrt{2}} |V_{ub}V_{ud}^*| e^{-i\gamma} f_\pi m_B^2 f_+(0) (c_1 + \frac{c_2}{3}) , \quad (\text{A4})$$

and comparing it with (1) we get:

$$\frac{1}{2\sqrt{3}} A_2^f + \frac{1}{\sqrt{6}} A_0^f = (c_1 + \frac{c_2}{3}) . \quad (\text{A5})$$

From (A3) and (A5) we obtain:

$$A_0^f = \frac{\sqrt{2}}{3\sqrt{3}} (5c_1 - c_2) = 1.54 . \quad (\text{A6})$$

The amplitude of $\bar{B}_d \rightarrow \pi^0 \pi^0$ decay can be calculated analogously, or it can be constructed from A_0^f and A_2^f with the help of (2):

$$\begin{aligned} M_{\bar{B}_d \rightarrow \pi^0 \pi^0}^f &= \frac{G_F}{\sqrt{2}} |V_{ub} V_{ud}^*| e^{-i\gamma} f_\pi m_B^2 f_+(0) \left[-\frac{5c_1 - c_2}{9} + \frac{8c_1 + 8c_2}{9} \right] = \\ &= \frac{G_F}{\sqrt{2}} |V_{ub} V_{ud}^*| e^{-i\gamma} f_\pi m_B^2 f_+(0) (c_2 + \frac{c_1}{3}) , \end{aligned} \quad (\text{A7})$$

a well known prediction of smallness of $\bar{B}_d \rightarrow \pi^0 \pi^0$ decay probability in the factorization approximation. However if FSI phase shift between A_0 and A_2 is large, this compensation disappears.

QCD penguins generate an additional term to the weak interaction Hamiltonian, (A1):

$$\Delta \hat{H} = \frac{G_F}{\sqrt{2}} (-) V_{tb} V_{td}^* [c_3 O_3 + c_4 O_4 + c_5 O_5 + c_6 O_6] , \quad (\text{A8})$$

$$\begin{aligned} O_3 &= \bar{d} \gamma_\alpha (1 + \gamma_5) b [\bar{u} \gamma_\alpha (1 + \gamma_5) u + \bar{d} \gamma_\alpha (1 + \gamma_5) d] , & c_3 &= 0.013 , \\ O_4 &= \bar{d}_a \gamma_\alpha (1 + \gamma_5) b^c [\bar{u}_c \gamma_\alpha (1 + \gamma_5) u^a + \bar{d}_c \gamma_\alpha (1 + \gamma_5) d^a] , & c_4 &= -0.032 , \\ O_5 &= \bar{d} \gamma_\alpha (1 + \gamma_5) b [\bar{u} \gamma_\alpha (1 - \gamma_5) u + \bar{d} \gamma_\alpha (1 - \gamma_5) d] , & c_5 &= 0.009 , \\ O_6 &= \bar{d}_a \gamma_\alpha (1 + \gamma_5) b^c [\bar{u}_c \gamma_\alpha (1 - \gamma_5) u^a + \bar{d}_c \gamma_\alpha (1 - \gamma_5) d^a] , & c_6 &= -0.037 . \end{aligned}$$

In order to find the penguin contribution to $B \rightarrow \pi\pi$ decay amplitudes let us calculate the matrix element of (A8) between $\bar{B}_d$ and $\pi^+ \pi^-$ in the factorization approximation. For this purpose it is convenient to use Fierz transformations of γ -matrices rewriting $\Delta \hat{H}$ in the following form:

$$\begin{aligned} \Delta \hat{H} &= -\frac{G_F}{\sqrt{2}} V_{tb} V_{td}^* [a_4 \bar{u} \gamma_\alpha (1 + \gamma_5) b \bar{d} \gamma_\alpha (1 + \gamma_5) u - \\ &\quad - 2a_6 \bar{u} (1 + \gamma_5) b \bar{d} (1 - \gamma_5) u] , \end{aligned} \quad (\text{A9})$$

where $a_4 = c_4 + 1/3c_3 = -0.027$, $a_6 = c_6 + 1/3c_5 = -0.034$.

Calculating the matrix element:

$$\langle \pi^+ \pi^- | \Delta \hat{H} | \bar{B}_d \rangle = -\frac{G_F}{\sqrt{2}} |V_{tb} V_{td}^*| e^{i\beta} f_+(0) f_\pi m_B^2 \left[a_4 + \frac{2m_\pi^2 a_6}{(m_u + m_d)(m_b - m_u)} \right] , \quad (\text{A10})$$

and comparing this expression with (1) we get:

$$P^f = -a_4 - \frac{2m_\pi^2}{(m_u + m_d)m_b} a_6 = +0.06 \pm 0.01 , \quad (\text{A11})$$

where $m_u + m_d = 9 \pm 3$ MeV, $m_b = 4.5$ GeV were substituted.

References

- [1] M.Gronau and D.London, Phys. Rev. Lett. **65**, 3381 (1990).
- [2] G.G. Ovanesyan and M.I.Vysotsky, Pisma v ZhETPh **81**, 449 (2005); JETP Letters **81**, 361 (2005).
- [3] R. Aleksan, F. Buccella, A. Le Yaouanc, L. Oliver, O. Pene, J.C. Raynal, Phys. Lett. **B356**, 95 (1995).
- [4] M.I.Vysotsky, hep-ph/0506322, Yad. Fiz. **69**, 703 (2006).
- [5] M. Bander, D. Silverman and A. Soni, Phys. Rev. Lett. **43**, 242 (1979);
G.M. Gérard and W.-S. Hou, Phys. Rev. **D43**, 2909 (1991).
- [6] M. Gronau, Phys. Rev. Lett. **63**, 1451 (1989).
- [7] HFAG, <http://www.slac.stanford.edu/xorg/hfag/>.
- [8] BABAR Collaboration, B. Aubert et al., Phys. Rev. **D72**, 051102 (2005).
- [9] A. Khodjamirian et al., Phys. Rev. **D72**, 094012 (2005).
- [10] E. Chell, M.G. Olsson, Phys. Rev. **D48**, 4076 (1993).
- [11] M. Shifman, A. Vainshtein, V. Zakharov, Zh. Eksp. Teor. Fiz. **72**, 1275 (1977);
A.I. Vainstein, Int. J. Mod. Phys. **A14**, 4705 (1999).
- [12] G.Buchalla, A.Buras, M.Lautenbacher, Rev. Mod. Phys. **68**, 1125 (1996).
- [13] Belle Colaboration, L.Widhalm et al., hep-ex/0604049 (2006).
- [14] CLEO Collaboration, P. Rubin et al., hep-ex/0512063 (2005);
CLEO Collaboration, M. Selen et al., Phys. Rev. Lett. **71**, 1973 (1993).
- [15] S. Eidelman et al., Phys. Lett. **B592**, 1 (2004) and update available on the PDG [www](http://www.pdg.lbl.gov/) pages.
- [16] CLEO Collaboration, S. Ahmed et al., Phys. Rev. **D66**, 031101 (2002).
- [17] A.Kaidalov, Proceedings of 24 Rencontre de Moriond “New results in hadronic interactions”, 391 (1989).
- [18] H.Y. Cheng, C.K. Chua and A. Soni, Phys. Rev. **D71**, 014030 (2005).

- [19] A.Deandrea et al., hep-ph/0508083 (2005).
- [20] J.P. Donoghue, E.Golowich, A.A. Petrov and J.M. Soares, Phys. Rev. Lett. **77**, 2178 (1996).
- [21] M.Suzuki, L.Wolfenstein, Phys. Rev. **D60**, 074019 (1999).
- [22] E.Alvarez, J.Pestieau, hep-ph/0602007 (2006).
- [23] K.G.Boreskov, A.A.Grigoryan, A.B.Kaidalov, I.I.Levintov, Yad. Fiz. **27**, 813 (1978).
- [24] M.Gronau, J.Rosner, Phys. Lett. **B595**, 339 (2004).
- [25] M.Hazumi, Plenary talk at ICHEP 06, Moscow, 2006.
- [26] BABAR Collaboration, B. Aubert et al., Phys. Rev. Lett **95**, 041805 (2005).
- [27] Belle Colaboration, K.Abe et al., hep-ex/0507039.
- [28] CKMfitter, <http://ckmfitter.in2p3.fr>.
- [29] UTfit, <http://utfit.roma1.infn.it>.